StandardBar[®]–Berry

Essential Vitamins, Minerals, and Other Important Nutrients in a Tasty but Healthy Snack

Who can we believe? Researchers, nutritionists, doctors, and other opinion leaders debate endlessly about the effects of eating too much or too little protein, too high or too low of a fat content, and what types and frequencies of exercise make a healthy difference in our lives. But in the midst of all of this dietary confusion, there are a few things that most experts agree upon: that most of us make too many unhealthy choices when it comes to snacks and meal planning. The recurrent theme that echoes throughout this debate is that of balance and nature when choosing what belongs in a healthy diet. Eating natural foods in proper balance equips the body with the fuel it requires to operate at peak efficiency. The 40-30-30 theory of dietary balance requires that we eat 40 percent of our daily caloric intake in carbohydrates, 30 percent in proteins, and 30 percent in fats in order to maintain balance.[†]

How Berry StandardBars Keep You Healthy

Supports cellular health

The blueberries, cranberries, cherries, and extra-virgin olive oil found in Berry StandardBars contain phytonutrients that are high in antioxidant activity.[†]

Promotes urinary and gastrointestinal tract health

Different phytonutrients from blueberries and cranberries help cleanse the urinary tract. Apple pectin, from apple fiber, helps maintain healthy intestinal flora.[†]

Maintains cardiovascular health

The flavonoids from cranberries help keep the arteries clear and healthy. Apples promote healthy lipid metabolism. Oats help keep the heart and blood vessels healthy. Calcium helps maintain a healthy heart. Black currant seed oil contains linoleic and gamma linolenic acids to help keep vascular walls healthy.[†]

Supports skeletal and immune system strength

Whey protein and calcium lactate provide a highly bioavailable source of calcium to build and maintain strong bones. Calcium, magnesium, and potassium work together to promote healthy muscle contraction. Black currant seed oil boosts the immune system.[†]



Introduced in 1999 GP 🚺

Content: Eighteen 1.75 oz. (50 g) bars

Supplement Facts: Serving Size: 1 bar Servings per Container: 18

	Amount	
	per Serving	%DV
Calories	195	
Calories from Fat	60	
Total Fat	7 g	11%*
Saturated Fat	0 g	0%*
Polyunsaturated Fat	1 g	
Monounsaturated Fat	3.5 g	
Cholesterol	0 mg	0%
Total Carbohydrate	23 g	7%*
Dietary Fiber	2 g	8%*
Sugars	16 g	
Protein	10 g	20%*
Calcium	100 mg	10%
Magnesium	28 mg	6%
Sodium	190 mg	8%
*Percent Daily Values (D	M are based on	

Amount

Percent Daily Values (DV) are based on a 2.000-calorie diet.

Ingredients: Brown rice syrup, whey (milk) protein, almond butter, pure wildflower honey, cherries, soybean lecithin, cranberries, apple juice concentrate, blueberries, glycerin, sesame seeds, cranberry puree, extra-virgin olive oil, black currant (seed) oil, blueberry puree, calcium lactate, oat fiber, apple fiber, and magnesium citrate.

Special Information: Free of cholesterol and saturated fat.

Sold through health care professionals.

Whole Food Philosophy

Our founder, Dr. Royal Lee, challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature-in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists-known and unknown-bioactivity is markedly enhanced over isolated nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to an isolated or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.



Please copy for your patients.

🕀 This product contains less than 10 parts per million of gluten per serving size or less than 20 parts per million per the suggested use listed on each product label. 🖤 Vegetarian (Lacto-ovo) +These statements have not been evaluated by the Food & Drug Administration. These products are not intended to diagnose, treat, cure, or prevent any disease

StandardBar[®]–Berry

What Makes Berry StandardBars Unique

Product Attributes

Ingredients are derived from whole food sources

- > A nutritious blend of three natural fruits: blueberries, cranberries, and cherries
- > Ingredients provide a nutritious snack alternative that supplies the body with many health-giving benefits
- > Whole food sources enhance the natural assimilation of nutrients and strength of synergistic cofactors, for example, the sesame seeds enhance tocopherol bioactivity
- > Combines healthy ingredients that support multiple organ systems
- Offers a convenient snack without artificial preservatives, colors, or flavors

Provides a healthy balance of carbohydrates, proteins, and fats

- > A 40-30-30 type health bar
- > Free of cholesterol and saturated fat
- > Enlists the properties of fruits and other nutrients well recognized for their protective and cleansing characteristics

Manufacturing and Quality-Control Processes Not disassociated into isolated components

> The nutrients in Berry StandardBar are processed to remain intact, complete nutritional compounds

Degreed microbiologists and chemists in our on-site laboratories continually conduct bacterial and analytical tests on raw materials, product batches, and finished products

> Ensures consistent quality and safety

Vitamin and mineral analyses validate product content and specifications

> Assures high-quality essential nutrients are delivered

Studies on nutrients generally use large doses and these studies, some of which are cited below, are the basis for much of the information we provide you in this publication about whole food medients. See the supplement facts for Berry StandardBar[®].

- Aprikian O., et al. 2002, Lyophilized apple counteracts the development of
- Aprikan O., et al. 2002. Lyopinized apple counteracts the development of hypercholestensina, oxidative stress, and rerail dysfunction in obese Zucker rats. Journal of Nutrition 132(7): 1969-1976. Bach J.F., Bach P.A. 1997. Prescription for Nutritional Healing. 2nd ed. Garden Clty Park, WY. Avery Publishing Group: 12-19, 23-27. Bertmeis M.C., et al. 1990. Selective effects of dietary tato on vascular 13-HODE synthesis and platiet/vessel wall interactions. Thrombosis Beasauch 50(16): 810.6-30.
- Research 59(5): 819-830. Bickford P.C., et al. 2000. Antioxidant-rich diets improve cerebral physiology
- and motor learning in aged rats. *Brain Research* 666(1-2): 211-217. Cooney R.V., et al. 2001. Effects of dietary sesame seeds on plasma tocopherol levels. *Nutrition and Cancer* 39(1): 66-71. Davy B.M., et al. 2002. High-fiber cat cereal compared with wheat cereal
- Davy B.M., et al. 2002. High-fiber cat correal compared with wheat cereal consumption favorably alters LDL-cholesterol subclass and particle numbers in middle-aged and older men. American Journal of Clinical Nutrition 76(2): 351-358. Deferns J.L., Leads A.R. 1996. Resting blood pressure and cardiovascular reactivity to mental antimetic in mild hypertensive males supplemented with black currant seed oil. Journal of Human Hypertension. Aug; 10(8): 531-537. Grong H, et al. 1998. Preventive and therapeutic effects of calcium lactate on estencomesis in and cardiochomized rats. Wei Shana Yan. Jin.
- on osteoporosis in aged ovariectomized rats. Wei Sheng Yan Jiu 27(6): 402-404. Hakkinen S.H., et al. 1999. Content of the flavonols quercetin, myricetin
- and kaempferol in 25 edible berries. Journal of Agrigulture and Food
- and kaempterol in 25 edulte bernes. Journal of Agryuiture and Foo Chemistry 47(6): 2274-2279. Ikeda S., et al. 2001. Dietary sesame seeds elevate alpha- and gamma-tocotrienio concentrations in sika and adjouse tissue of rafs fed the tootrienio-rich fraction extracted from paim oil. Journal of Nutrition 131(11): 2892-2897. Jenkins A.L., et al. 2002. Degression of the glycemic index by high levels of halta-ultera flops in two functional forsit best in two 2 (fabrates
- of beta-glucan fiber in two functional foods tested in type 2 diabetes. European Journal of Clinical Nutrition 56(7): 622-628. Kalt W., et al. 2001, Interspecific variation in anthocyanins, pheolic

- Katt W., et al. 2001. Interspecific variation in anthocyanins, pheolics, and antioxidant capacity among genotypes of highlbush and dowbush blueberriss (*Vaccinium section cyanocoaccus spp.*). *Journal of Agriculture and Food Chemistry* 49(10): 4761-4767. Keeran J.M., et al. 2002. Dati ingestion reduces systolic and diastolic blood pressure in patients with mild or borderline hypertension: a pilot trial. *Journal of Family Practice* 51(4): 369. Kris: Elmerton P.M., et al. 2002. High-soluble-fiber foods in conjunction with a telephone-tased, personalized behavior change support service result in favorable changes in lipids and lifestyles after 7 weeks. *Journal of the American Diabetes Association* 102(4): 503-510. Markus C.R., et al. 2002. Whey protein rich in alpha-factationmin increases the ratio of plasma tryptophen to the sum of the other large neutral amino
- the ratio of plasma tryptophan to the sum of the other large neutral amino acids and improves cognitive performance in stress-vulnerable subjects. American Journal of Clinical Nutrition 75(6): 1051-1056.
- American Journal or Cunical Nutrition 7 s(b): 1051-1056.
 Micke P., et al. 2002. Effects of long-term supplementation with whey proteins on plasma glutathione levels of HIV-infected patients. *European Journal of Nutrition* 41(1): 12-18.
 Katarajan S., et al. 2001. Healing of an MRSA-colonized, hydroxyurea-induced leg ulcer with honey. *Journal of Dermatology Treatment* 17(1): 33-96.
- Pan Z., et al. 2000. Effects of oral calcium supplementation on blood
- pressure in population. *Zhonghua Yu Fang Yi Xue Za Zhi* 34(2): 109-112. Pins J.J., et al. 2002. Do whole-grain oat cereals reduce the need for
- antihypertensive medications and improve blood pressure control? Journal of Family Practice 51(4): 353-359.
- Pitchford P. 1993. *Healing With Whole Foods*. Revised ed. Berkeley, CA: North Atlantic Books: 144-145, 151, 178-179, 288, 317, 319, 429, 470, 492, 576-578.
- 410, 432, JPC-31, Horn Librard, Alexandrian, Alexandrian, Alexandria, Alexa
- Reed J. 2002. Cranberry flavonoids, atherosclerosis and cardiovascular health. Critical Review of Food Science Nutrition 43(3 Suppl): 301-316
- Reid G. 2002. The role of cranberry and probiotics in intestinal and urogenital tract health. *Critical Review of Food Science Nutrition* 42(3 Suppl): 293-300
- Reid G., et al. 2001. Cranberry juice consumption may reduce biofilms on uroepithelial cells: pilot study in spinal cord injured patients. Spinal Cord
- uroepithelial cells: pilot study in spinal cord injured patients. Spinal Cord 39(1): 26-30. Roffe C., et al. 2002. Randomised, cross-over, placebo controlled trial of magnesium critate in the treatment of chronic persistent leg cramps. *Medical Science Monitor* 8(5): CR326-CR330. Sattmann E, et al. 2001. An cat-containing hypocaloric diet reduces systolic blood pressure and improves lingl profile beyond offects of weight loss in man producemen. *Journal of Multition* 33(16): 1465-1470.
- men and women. Journal of Nutrition 131(5): 1465-1470.
- Seeram N.P., et al. 2001. Cyclooxygenase inhibitory and antioxidant cyaniding glycosides in cherries and berries. *Phytomedicine* 8(5): 362-369
- 382-389.
 Seligan S., et al. 2002. Phenolic compounds and antioxidant capacity of Georgia-grown blueberries and blackberries. *Journal of Agriculture and Food Chemistry 50(8): 2432-2438.*Strato-Yasumolo S., et al. 2001. Effect of sesame seeds rich in sesamin and sesamilin on fatty acid oxidation in rat liver. *Journal of Agriculture and Food Chemistry* 49(5): 2647-2651.
 Spiller G.A., et al. 1998. Nuts and plasma lipids: an almond-based diet lowers LDL-C while preserving HDLC. *Journal of the American College of Nutrition* 17(3): 285-290.

- College of Nutrition 17(3): Zeo-290. Takeuchi H., et al. 2001. Hypoglycemic effect of a hot-water extract from defatted sesame (sesamum indicum L.) seed on the blood glucose level in genetically diabetic KK-Ay mice. *Bioscience, Biotechnology, and*
- in genetically diabetic KK-Ay mice. *Bioscience, Biotechnology, and Biochemistry* 66(10): 2318-2321. Tazawa K, et al. 1999. Dietary fiber inhibits the incidence of hepatic metastasis with the anti-oxidant activity and portal scavenging functions. *Human Cell* 12(4): 189-196. Truswell A.S. 2002. Cereal grains and coronary heart disease. *European Journal of Clinical Nutrition* 56(1): 1-14. *Vieis IE*, 6alf C. 2002. Biochemical companying and the other hemicals
- Visiol F., Galli C. 2002. Biological properties of olive oil phytochemicals.
- Critical Review of Food Science and Nutrition 42(3): 209-221 Wu D., et al. 1999. Effect of dietary supplementation with black currant
- seed oil on the immune response of healthy elderly subjects. American Journal of Clinical Nutrition 70(4): 536-543. Youdim K.A., et al. 2000. Polyphenolics enhance red blood cell resistance to oxidative stress: in vitro and in vivo. Biochem Biophys Acta 1523(1): 117-122.

