Cruciferous Complete[™]

Contains Organically Grown, Minimally Processed Kale and Brussels Sprouts

Research demonstrates that cruciferous vegetables, such as kale and Brussels sprouts, contain important phytonutrients that help protect against free radicals, the highly unstable molecules that can damage cells and genetic material. Cruciferous vegetables are nutrient-dense foods packed with many essential vitamins, minerals, and antioxidants. Cruciferous vegetables also contain important health-promoting nutrients like carotenoids and other phytochemicals that stimulate enzymatic activity to support liver detoxification.[†]

How Cruciferous Complete Keeps You Healthy

Furnishes many nutrients to support health

Cruciferous Complete contains kale and Brussels sprouts, which contain a broad spectrum of micronutrients, including vitamins B_6 , C, and K; dietary fiber; copper; calcium; and potassium. A class of plant pigments, called carotenoids, is also found in crucifers. These pigments, including beta carotene and lutein, serve important roles within the body. Beta carotene is a precursor to vitamin A, while lutein is found in abundance in the retina of the eye. Carotenoids and other phytochemicals also have antioxidant properties and can protect cells from damage of excess reactive oxygen species. †

Provides antioxidant support

The health-promoting phytochemicals in cruciferous vegetables protect cells against free-radical damage and have been shown in the laboratory to alter the metabolic pathways responsible for the metabolism of hormones in cultured cells and in mice.

Induces liver detoxification enzymes

Cruciferous vegetables contain highly active phytochemicals called glucosinolates. In laboratory cell cultures and mice, products left from the breakdown of glucosinolates stimulate the phase I and phase II liver enzymes, two biochemical pathways in the liver involved in converting toxins into harmless substances that can easily be excreted from the body.



Introduced in 1998

Œ

Content: 90 capsules

Suggested Use: One capsule per day, or as directed.

Supplement Facts: Serving Size: 1 capsule Servings per Container: 90

 Amount per Serving
 %DV

 Calories
 3

 Vitamin K
 4 mcg
 4%

 Potassium
 10 mg
 <1%</td>

 Kale
 300 mg

 Brussels sprouts
 300 mg

Ingredients: See Supplement Facts.

Other Ingredients: Gelatin, water, calcium stearate, and colors.

Sold through health care professionals.

Please copy for your patients.



Cruciferous Complete[™]

What Makes Cruciferous Complete Unique

Product Attributes

Ingredients are derived from whole food sources

Each capsule supplies 300 mg each of organically grown kale and Brussels sprouts

Certified Organic Farming

A healthy ecosystem is created by using organic farming techniques, such as rotating crops, fertilizing the soil with nutrient-rich cover crops and byproducts from our processing, practicing strict weed-control standards, and continually monitoring the health of our plants

- > Assures the soil is laden with minerals and nutrients
- Ensures plants are nutritionally complete and free from synthetic pesticides

Unique Processing

Upon harvesting, nutrient-rich plants are immediately washed and promptly processed

> Preserves nutritional integrity

Low-temperature, high-vacuum drying technique

> Preserves the enzymatic vitality and nutritional potential of ingredients

Not disassociated into isolated components

> The nutrients in Cruciferous Complete are processed to remain intact, complete nutritional compounds

Degreed microbiologists and chemists in our on-site laboratories continually conduct bacterial and analytical tests on raw materials, product batches, and finished products

> Ensures consistent quality and safety

Vitamin and mineral analyses validate product content and specifications

> Assures high-quality essential nutrients are delivered

Whole Food Philosophy

Our founder, Dr. Royal Lee, challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature—in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists-known and unknown—bioactivity is markedly enhanced over isolated nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to an isolated or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.

Studies on nutrients generally use large doses and these studies, some of which are cited below, are the basis for much of the information we provide you in this publication about whole food ingredients. See the supplement facts for Cruciferous Complete $^{\rm TM}$.

Auborn, K. J., Fan, S., Rosen, E. M., Goodwin, L., Chandraskaren, A. Williams, D. E., et al. (2003). Indole-3-carbinol is a negative regulator of estrogen. J Nutr, 133(7 Suppl), 2470S-2475S.
Hecht, S. S. (1999). Chemoprevention of Cancer by Isothiocyanates,

Modifiers of Carcinogen Metabolism, J. Nutr., 129(3), 768.

Hecht, S. S. (2000). Inhibition of carcinogenesis by isothiocyanates. *Drug Metab Rev*, 32(3-4), 395-411.

Holick, C. N., Michaud, D. S., Stolzenberg-Solomon, R., Mayne, S. T.,

Pietinen, P., Taylor, P. R., et al. (2002). Dietary carotenoids, serum beta-carotene, and retinol and risk of lung cancer in the alpha-tocopherol, beta-carotene cohort study. Am J Epidemiol, 156(6), 536-547. Holst, B., Williamson, G. (2004), A critical review of the bioavailability of

glucosinolates and related compounds. Nat Prod Rep., 21(3), 425-447. Institute of Medicine Food and Nutrition Board. (2000). Beta-carotene and other carotenoids. In Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids (pp. 325-400). Washington D.C.:

International Agency for Research on Cancer. (1998). IARC Handbooks of Cancer Prevention: Carotenoids, Lyon: International Agency for Research on Cancer.

Nessell of Ortalicel.

Johnston, N. (2004). Sulforaphane halls breast cancer cell growth. *Drug Discovery Today*, 9(21), 908.

Keck, A.-S., & Finley, J. W. (2004). Cruciferous Vegetables: Cancer

Protective Mechanisms of Glucosinolate Hydrolysis Products and Selenium. *Integr Cancer Ther*, 3(1), 5-12.
Krinsky, N. I., Landrum, J. T., & Bone, R. A. (2003). Biologic mechanisms

of the protective role of lutein and zeaxanthin in the eve. Annu Rev Nutr.

23, 171-201.

Kristal, A., Lampe JW. (2002). Brassica vegetables and prostate cancer risk: a review of the epidemiological evidence, Nutr Cancer, 42(1), 1-9,

Liu, R. H. (2004). Potential Synergy of Phytochemicals in Cancer Prevention: Mechanism of Action. J. Nutr., 134(12), 3479S-3485.
Mares-Perlman, J. A., Millen, A. E., Ficek, T. L., & Hankinson, S. E. (2002). The body of evidence to support a protective role for lutein and zeaxanthin

in delaying chronic disease. Overview. *J Nutr*, 132(3), 518S-524S. Munday, R., & Munday, C. M. (2004). Induction of phase II detoxification enzymes in rats by plant-derived isothiocyanates: comparison of ally isothiocyanate with sulforaphane and related compounds. J Agric Food

Chem, 52(7), 1867-1871.
Olson, J. A. (1999). Carotenoids and human health. Arch Latinoam Nutr, 49(3 Suppl 1), 7S-11S.

Rao, A. V., & Rao, L. G. (2007). Carotenoids and human health. *Pharmacol*

Res, 55(3), 207-216.
Telang, N. T., Katdare, M., Bradlow, H. L., Osborne, M. P., & Fishman, J (1997), Inhibition of proliferation and modulation of estradiol metabolism (1997), Illimitation in professional and includation to seasonal miscardinal novel mechanisms for breast cancer prevention by the phytochemical indole-3-carbinol. Proc Soc Exp Biol Med, 216(2), 246-252. van Poppel, G., Verhoeven, D. T., Verhagen, H., & Goldbohm, R. A.

(1999). Brassica vegetables and cancer prevention. Epidemiology and mechanisms. Adv Exp Med Biol, 472, 159-168. Verhoeven, D. T., Goldbohm, R. A., van Poppel, G., Verhagen, H., van den Brandt, P. A. (1996). Epidemiological studies on Brassica vegetables and

cancer risk. Cancer Epidemiol Biomarkers Prev, (5(9), 733-748.
Walters, D. G., Young, P. J., Agus, C., Krite, M. G., Boobis, A. R.,
Gooderham, N. J., et al. (2004). Cruciferous vegetable consumption
alters the metabolism of the dietary carcinogen 2-amino-1-methyl-

alters the metabolism of the delary carcinogen 2-amino-1-methyl-e-phenylmidzol/4,5-b] pyridine (PhiP) in humans. Carcinogenesis, 25(9), 1659-1669.

Young, A. J., & Lowe, G. M. (2001). Antioxidant and prooxidant properties of carotenoids. Arch Biochem Biophys, 385(1), 20-27.

Zhang, Y. (2004). Cancer-preventive isothicoyanates: measurement of human exposure and mechanism of action. Mutat Res, 555(1-2), 173, 190.

Ziegler, R. G. (1989). A review of epidemiologic evidence that carotenoids reduce the risk of cancer. *J Nutr*, 119(1), 116-122.

